

organs) the ants did not respond aggressively.

Not only were the behaviors of the attendant ants changed but also dopamine levels in their brains, which increased significantly in ants that attended caterpillars with functioning nectary organs. Other biogenic amines (serotonin, octopamine and tyramine) were not affected. Exactly how the secretions from the nectary organs could elicit such changes in an important neuromodulator and neurotransmitter remains unknown. The quantity of secretions from the nectary organs is very small, making a complete chemical characterization difficult. In addition, the researchers only measured four biogenic amines, but other changes in the brains of attendant ants might be taking place. Hopefully, future work such as whole brain metabolomics or RNA-seq experiments on individual brains will provide further insights.

But the importance of this paper is not just for the details it provides on the proximate mechanisms of behavioral manipulation [11]. The importance is also that apparently mutualistic caterpillars manipulate ant behavior at all. It is well known that parasites can adaptively manipulate the behavior of their hosts. This is the concept of the extended phenotype [12], where changes in host behavior benefit parasite fitness. In the

study by Hojo *et al.* [2], it seems that attending bodyguard ants are less likely to wander from their charge and more likely to be aggressive, which should benefit the caterpillar. This ‘mutualism’ thus has all the hallmarks of adaptive manipulation of host behavior by a parasite!

It would appear that caterpillars enforce the cooperation they require. This is likely to be due to the fact that the ant colony may not need its caterpillar ‘sugar tap’ as much as the sugar tap needs its fierce bodyguards. As other sources of sugar present themselves, the danger for the caterpillar is that the ants shift away from their protective role, leaving the caterpillar vulnerable to predation. And so, perhaps by way of an insurance mechanism, the hungry caterpillar has evolved to keep their ant bodyguards on a short leash using manipulative drugs. This study will hopefully encourage different researchers to examine other apparent manipulations for signs of similar manipulative behaviors.

## REFERENCES

1. Rico-Gray, V., and Oliveira, P.S. (2007). *The Ecology and Evolution of Ant-Plant Interactions* (Chicago: University of Chicago Press).
2. Hojo, M.K., Pierce, N.E., and Tsuji, K. (2015). Lycaenid caterpillar secretions manipulate attendant ant behavior. *Curr. Biol.* 25, 2260–2264.
3. Hölldobler, B., and Wilson, E.O. (2009). *The superorganism: the beauty, elegance, and strangeness of insect societies* (W.W. Norton).
4. Hölldobler, B., and Wilson, E.O. (1990). *The Ants* (Cambridge, Mass.: Harvard University Press).
5. Davidson, D.W., Cook, S.C., Snelling, R.R., and Chua, T.H. (2003). Explaining the abundance of ants in lowland tropical rainforest canopies. *Science* 300, 969–972.
6. Pierce, N.E., and Mead, P.S. (1981). Parasitoids as selective agents in the symbiosis between lycaenid butterfly larvae and ants. *Science* 211, 1185–1187.
7. Pierce, N.E. (1995). Predatory and parasitic Lepidoptera: carnivores living on plants. *J. Lepid. Soc.* 49, 412–453.
8. Als, T.D., Vila, R., Kandul, N.P., Nash, D.R., Yen, S.H., Hsu, Y.F., Mignault, A.A., Boomsma, J.J., and Pierce, N.E. (2004). The evolution of alternative parasitic life histories in large blue butterflies. *Nature* 432, 386–390.
9. Pierce, N.E., Braby, M.F., Heath, A., Lohman, D.J., Mathew, J., Rand, D.B., and Travassos, M.A. (2002). The ecology and evolution of ant association in the Lycaenidae (Lepidoptera). *Annu. Rev. Entomol.* 47, 733–771.
10. Hojo, M.K., Yamamoto, A., Akino, T., Tsuji, K., and Yamaoka, R. (2014). Ants use partner specific odors to learn to recognize a mutualistic partner. *PLoS One* 9, e86054.
11. Hughes, D.P. (2013). Pathways to understanding the extended phenotype of parasites in their hosts. *J. Exp. Biol.* 216, 142–147.
12. Dawkins, R. (1982). *The extended phenotype* (Oxford: W.H. Freeman).

# Plant Intracellular Transport: Tracing Functions of the Retrograde Kinesin

Sabine Müller

ZMPB, University of Tübingen, Morganstelle 32, 72026, Tübingen, Germany

Correspondence: [sabine.mueller@zmbp.uni-tuebingen.de](mailto:sabine.mueller@zmbp.uni-tuebingen.de)

<http://dx.doi.org/10.1016/j.cub.2015.08.011>

**Adding to its varied repertoire of functions in cell morphogenesis and cell division, a molecular motor protein of the kinesin-14 class has recently been implicated in rapid retrograde transport along cellular tracks in moss.**

Like traffic in general, cellular trafficking also relies on dedicated tracks and vehicles that facilitate the transport of cargo. In all eukaryotes, transport tracks

are made of the cytoskeleton — polar microtubules and actin filaments. Aided by versatile molecular motors, the list of cellular mechanisms requiring the

cytoskeleton is nearly infinite, and includes diverse functions in cell division, cell polarization and growth as well as in subcellular transport. In most eukaryotes,

cargo transport along these cellular tracks is assisted by three motor protein families — the microtubule-dependent kinesins, the dyneins and the actin-dependent myosins. However, the microtubule minus-end directed dyneins were lost from flowering plants and are absent from most clades of the lower plants [1], raising the question of exactly how plant cells accomplish minus-end directed, retrograde microtubule-dependent transport. Members of the minus-end directed kinesin-14 class motor proteins might substitute for dynein to achieve retrograde transport functions; yet, until recently, processive motility was only confirmed for kinesin-14 in yeast, but was reported neither for animal nor plant kinesin-14.

In a new study, published recently in *Nature Plants*, [2], Jönsson and co-workers now show that a member of the kinesin-14 class motor proteins in the moss *Physcomitrella patens* performs retrograde transport towards the minus-end of microtubules. In moss, the kinesin-14 class encompasses six subdivisions, kin-14I to kin-14VI, and based on subcellular localization they were assigned mitotic and non-mitotic functions [3]. The authors used advanced imaging of *in vitro* microtubule gliding assays to visualize the directional displacement of polarity-marked microtubules by purified kinesin-14 motors attached to glass cover slips. Four of the six representative kinesin-14s tested displayed microtubule minus-end directed motility as dimers [2]. To analyze the motors' processivity *in vitro*, single-molecule motility assays were performed, where labeled microtubules were bound to a cover slip and dilute kinesin-14 preparations were added [2]. Processive motors stay attached to the microtubule, even when walking over long distances, while non-processive motors detach immediately after each step. In these experiments, none of the investigated kin-14 motor-dimers demonstrated processive movement [2].

Nevertheless, one motor domain, kin-14Vlb, displayed particularly fast microtubule gliding velocity and, therefore, the full-length kin14Vlb (FL) protein was examined for processivity. Despite the majority of dimeric kin-14Vlb (FL) remaining non-processive, some brighter, mobile particles revealed the

presence of a small number of processively moving kin-14Vlb. This processive movement originated from the clustering of two or more kin-14Vlb dimers, akin to the processive movement of an artificial kin-14Vlb tetramer [2]. In support of the notion that the processive behavior of kin-14Vlb depended on multiple motor dimers, clustering of kin-14Vlb on liposomes enabled liposome transport along microtubules [2].

Finally, fluorescently labeled, endogenous kin-14Vlb punctae moved processively along microtubules in moss protonemal cells, and comparison of signal intensities with protein complexes of known fluorophore numbers suggested that the mobile kin-14Vlb punctae indeed contained more than one dimer [2]. The authors conclude that the clustering of a few non-processive kin-14Vlb dimers collectively enables the processive microtubule minus-end directed movement in the moss. Although the potential cargos are still elusive, it seems that the search for the retrograde microtubule-dependent motor has come to a closure in the moss.

However, this finding leads to the question of which functions kin-14Vlb motor might have in other plants. Indeed, the homologue of kin-14Vlb in flowering plants, kinesin-like calmodulin-binding protein (KCBP), has been extensively studied *in vitro* and *in vivo* [4–9]. KCBP binding to calcium-calmodulin inhibits the motor's microtubule binding ability [10–12] and likely this is also the case for moss kin-14Vlb. *Arabidopsis* KCBP is a minus-end directed motor *in vitro* [10], and in cotton KCBP displays punctate localization along microtubules [7] in agreement with the findings for moss kin-14Vlb [2]. Further indicative of a possible transport function of KCBP is the formation of abnormal leaf hairs in the *Arabidopsis* KCBP mutant *zwichel* [5]. Yet processive motility, as shown for moss kin-14Vlb, has not been reported for flowering plant KCBP so far. In addition to the non-mitotic function, the flowering plant KCBPs serve diverse roles in cell division — KCBPs associate with mitotic microtubule arrays and inhibition of KCBP function during mitosis delays cytokinesis [4,7–9], features that have not been reported for moss kin-14Vlb [2,3].

Recently, adding a new twist to KCBP's function in flowering plants, KCBP was

attributed to the plant specific mode of division plane positioning [8]. Plant cells are immobilized within tissues due to their confinement by cell walls; therefore, spatio-temporal control over cell divisions is crucial for plant development. Plant cells do not divide by constriction, but instead form a new partitioning wall, the cell plate, in the cell center during cytokinesis. KCBP was recently implicated in the guidance mechanism that attracts the cell plate to its final destination [8], which is pre-selected prior to mitosis by a transient band of microtubules, the plant-specific preprophase band [13]. The preprophase band disassembles, but leaves behind a set of 'landmark' proteins, collectively referred to as the cortical division site, that is required to preserve the positional information of the preprophase band throughout mitosis [13,14]. The recently revealed presence of KCBP at the cortical division site paves the way for speculation whether KCBP-mediated forces might contribute to the unique cell plate guidance mechanisms [8].

Thus, the kin-14Vlb/KCBP microtubule-minus end-directed motors emerge as versatile motor proteins that replace dynein as a retrograde transporter in the moss, and likely in the entire plant lineage, pending proof of processive movement. It should be noted here that rapid subcellular transport has been attributed to the actin-myosin cytoskeleton [15]. However, plant-specific mitotic KCBP functions might have evolved concurrent with the transition from simple to complex plant architecture in the context of cell wall-imposed positional confinement. Consistent with their plant-specific functions, the kin-14Vlb/KCBPs are found only in plants and in green algae, underpinning their evolution in the green lineage [16,17]; their motor domain, however, originates before the divergence of plants and animals [17].

## REFERENCES

1. Lawrence, C.J., Morris, N.R., Meagher, R.B., and Dawe, R.K. (2001). Dyneins have run their course in plant lineage. *Traffic* 2, 362–363.
2. Jonsson, E., Yamada, M., Vale, R.D., and Goshima, G. (2015). Clustering of a kinesin-14 motor enables processive retrograde microtubule-based transport in plants. *Nat. Plants* 1, 15087.

3. Miki, T., Naito, H., Nishina, M., and Goshima, G. (2014). Endogenous localizome identifies 43 mitotic kinesins in a plant cell. *Proc. Natl. Acad. Sci. USA* **111**, E1053–E1061.
4. Bowser, J., and Reddy, A.S. (1997). Localization of a kinesin-like calmodulin-binding protein in dividing cells of *Arabidopsis* and tobacco. *Plant J.* **12**, 1429–1437.
5. Oppenheimer, D.G., Pollock, M.A., Vacik, J., Szymanski, D.B., Ericson, B., Feldmann, K., and Marks, M.D. (1997). Essential role of a kinesin-like protein in *Arabidopsis* trichome morphogenesis. *Proc. Natl. Acad. Sci. USA* **94**, 6261–6266.
6. Smirnova, E.A., Reddy, A.S., Bowser, J., and Bajer, A.S. (1998). Minus end-directed kinesin-like motor protein, Kcbp, localizes to anaphase spindle poles in *Haemanthus* endosperm. *Cell Motil. Cytoskel.* **41**, 271–280.
7. Preuss, M.L., Delmer, D.P., and Liu, B. (2003). The cotton kinesin-like calmodulin-binding protein associates with cortical microtubules in cotton fibers. *Plant Physiol.* **132**, 154–160.
8. Buschmann, H., Dols, J., Kopischke, S., Peña, E.J., Andrade-Navarro, M.A., Heinlein, M., Szymanski, D.B., Zachgo, S., Doonan, J.H., and Lloyd, C.W. (2015). *Arabidopsis* KCBP interacts with AIR9 but stays in the cortical division zone throughout mitosis via its MyTH4-FERM domain. *J. Cell Sci.* **128**, 2033–2046.
9. Vos, J.W., Safadi, F., Reddy, A.S.N., and Hepler, P.K. (2000). The kinesin-like calmodulin binding protein is differentially involved in cell division. *Plant Cell* **12**, 979–990.
10. Song, H., Golovkin, M., Reddy, A.S., and Endow, S.A. (1997). In vitro motility of AtKCBP, a calmodulin-binding kinesin protein of *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **94**, 322–327.
11. Kao, Y.L., Deavours, B.E., Phelps, K.K., Walker, R.A., and Reddy, A.S. (2000). Bundling of microtubules by motor and tail domains of a kinesin-like calmodulin-binding protein from *Arabidopsis*: regulation by Ca(2+)/Calmodulin. *Biochem. Biophys. Res. Commun.* **267**, 201–207.
12. Narasimhulu, S.B., and Reddy, A.S.N. (1998). Characterization of microtubule binding domains in the *Arabidopsis* kinesin-like calmodulin binding protein. *Plant Cell* **10**, 957–965.
13. Rasmussen, C.G., Wright, A.J., and Müller, S. (2013). The role of the cytoskeleton and associated proteins in plant cell division plane determination. *Plant J.* **75**, 258–269.
14. Lipka, E., Gadeyne, A., Stöckle, D., Zimmermann, S., De Jaeger, G., Ehrhardt, D.W., Kirik, V., Van Damme, D., and Müller, S. (2014). The phragmoplast-orienting kinesin-12 class proteins translate the positional information of the preprophase band to establish the cortical division zone in *Arabidopsis thaliana*. *Plant Cell* **26**, 2617–2632.
15. Shimmen, T., and Yokota, E. (2004). Cytoplasmic streaming in plants. *Curr. Opin. Cell Biol.* **16**, 68–72.
16. Abdel-Ghany, S.E., Day, I.S., Simmons, M.P., Kugrens, P., and Reddy, A.S.N. (2005). Origin and evolution of kinesin-like calmodulin-binding protein. *Plant Physiol.* **138**, 1711–1722.
17. Dymek, E.E., Goduti, D., Kramer, T., and Smith, E.F. (2006). A kinesin-like calmodulin-binding protein in *Chlamydomonas*: evidence for a role in cell division and flagellar functions. *J. Cell Sci.* **119**, 3107–3116.